arXiv:1007.1729v1 [math.NT] 10 Jul 2010

# GALOIS GROUPS AND GENERA OF A KIND OF QUASI-CYCLOTOMIC FUNCTION FIELDS

MIN SHA AND LINSHENG YIN

Abstract. We call a $(q-1)$-th Kummer extension of a cyclotomic function field a quasi-cyclotomic function field if it is Galois, but non-abelian, over the rational function field with the constant field of $q$ elements. In this paper, we determine the structure of the Galois groups of a kind of quasi-cyclotomic function fields over the base field. We also give the genus formulae of them.

## 1. Introduction

We call a $(q-1)$-th Kummer extension of a cyclotomic function field a quasi-cyclotomic function field if it is Galois, but non-abelian, over the rational function field $k = \mathbb{F}_q(T)$. A large kind of such fields are described explicitly in [4] following the works in [1] and in [2]. In this paper, we describe the Galois groups of this kind of quasi-cyclotomic function fields by generators and relations following the method in [7] by using the results in [2] and [4]. We also give the genus formulae of them.

Now we recall the constructions of the quasi-cyclotomic function fields in [4]. Let $k = \mathbb{F}_q(T)$ be the rational function field over the finite field $\mathbb{F}_q$ of $q$ elements. In this paper we always assume the characteristic of $k$ is an odd prime number $p$. Put $\mathbb{A} = \mathbb{F}_q[T]$. Let $\Omega$ be the completion of the algebraic closure of the completion $\mathbb{F}_q((1/T))$ of $k$ at the place $1/T$. Let $k^{ac}$ be the algebraic closure of $k$ in $\Omega$. Let $k^{ab}$ be the maximal abelian extension of $k$ in $k^{ac}$.

Let $\bar{\pi} \in \Omega$ be the period of the Carlitz module, namely the lattice $\bar{\pi}\mathbb{A}$ of rank one corresponds to the Carlitz module, $\mathbf{e}_C$ the Carlitz exponential function defined by

$$\mathbf{e}_C(x) = x \prod_{0 \neq u \in \bar{\pi}\mathbb{A}} \big(1 - \frac{x}{u}\big), \quad x \in \Omega.$$

For $A \in \mathbb{F}_q((1/T))$, let $\{A\}$ be the representation in $\big(\mathbb{F}_q((1/T)) \setminus \mathbb{F}_q[T]\big) \cup \big\{0\big\}$ of $A$ modulo $\mathbb{A}$, we define

$$\sin(A) = \sqrt[q-1]{-1} \cdot \mathbf{e}_C(\bar{\pi}\{A\}/\mathrm{sgn}(\{A\})),$$

where sgn is the sign function on $\mathbb{F}_q((1/T))$.

Let $\mathcal{A}$ be the free ablian group generated by the symbols $[A] \in k/\mathbb{A}$. Define two homomorphisms

$$\mathbf{e}, \sin\colon \mathcal{A} \to k^{ab*}$$

---

2010 *Mathematics Subject Classification.* Primary 11R58; Secondary 11R32, 11R60.

*Key words and phrases.* quasi-cyclotomic function field, Galois group, genus.

This work was supported by NSFC Project No.10871107.

such that $\mathbf{e}([A]) = \mathbf{e}_C(\bar{\pi}A)$ and $\sin[A] = \sin(A)$ if $A \notin \mathbb{A}$, and $\mathbf{e}([A]) = 1$ and $\sin([A]) = 1$ otherwise.

Fix a total order $<$ in $\mathbb{A}$. Write $d_A$ for the degree of $A \in \mathbb{A}$. Let $M \in \mathbb{A}$ be monic. Put

$$S_M = \Big\{\text{monic prime factors of } M\Big\}.$$

Fix a generator $\gamma$ of $\mathbb{F}_q^*$. For $P, Q \in S_M$ with $P < Q$, let

$$\mathbf{a}_{PQ} = \sum_{\substack{d_A < d_Q \\ A:\text{monic}}} \sum_{\substack{d_B < d_P \\ B:\text{monic}}} \sum_{s=1}^{q-2} s\Big([\frac{BQ + \gamma^{-s}A}{PQ}] - [\frac{AP + \gamma^{-s}B}{PQ}]\Big).$$

It is easy to see $\sin\mathbf{a}_{PQ} = \mathbf{e}(\mathbf{a}_{PQ})$. We put

$$u_{PQ} = \begin{cases} \sin\mathbf{a}_{PQ}, & \text{if } 2|d_P, 2|d_Q \\ \\ \sqrt{P}\sin\mathbf{a}_{PQ}, & \text{if } 2|d_P, 2 \nmid d_Q \\ \\ \sqrt{Q}\sin\mathbf{a}_{PQ}, & \text{if } 2 \nmid d_P, 2|d_Q \\ \\ \sqrt{PQ}\sin\mathbf{a}_{PQ}, & \text{if } 2 \nmid d_P, 2 \nmid d_Q \end{cases}$$

Set $K = k(\mathbf{e}_C(\frac{\bar{\pi}}{M}))$, which is the cyclotomic function field of conductor $M$ and is abelian over $k$. For $P \in S_M$, write $P^* = (-1)^{\frac{|P|-1}{q-1}}P = (-1)^{d_P}P$, where $|P| = q^{d_P}$, then $\sqrt[q-1]{P^*} \in K$, hence $\sqrt{P^*} \in K$. As $\sin\mathbf{a}_{PQ} \in K$, we have $u_{PQ} \in K$. Put $\widetilde{K} = K(\sqrt[q-1]{u_{PQ}})$. By [Sect.5.2, 4], $\widetilde{K}$ is a quasi-cyclotomic function field over $k$, whose degree over $K$ equals to $q-1$.

## 2. The Galois groups

Let $G = \mathrm{Gal}(K/k)$ and $\widetilde{G} = \mathrm{Gal}(\widetilde{K}/k)$ be the Galois groups of the extensions $K/k$ and $\widetilde{K}/k$ respectively. In this section, we determine $\widetilde{G}$ by generators and relations.

Write $w = q-1$ and $u = u_{PQ}$ for simplicity. Clearly $\mathrm{Gal}(\widetilde{K}/K)$ is isomorphic to $\mathbb{Z}/w\mathbb{Z}$. Recall that $\gamma$ is a fixed generator of $\mathbb{F}_q^*$. Let $\epsilon \in \mathrm{Gal}(\widetilde{K}/K)$ be a generator such that $\epsilon(\sqrt[w]{u}) = \gamma\sqrt[w]{u}$. Denote the isomorphism $\mathbb{F}_q^* \to \mathbb{Z}/w\mathbb{Z}$, $\gamma^i \mapsto \bar{i}$ by $\log_\gamma$. Each element of $G$ has $w$ liftings in $\widetilde{K}$.

**Lemma 2.1.** *For $\sigma \in G$, choose $v_\sigma \in K^*$ such that $\sigma(u) = v_\sigma^w u$, we define a lifting $\tilde{\sigma} \in \widetilde{G}$ of $\sigma$ by $\tilde{\sigma}(\sqrt[w]{u}) = v_\sigma\sqrt[w]{u}$. Then $\widetilde{G} = \{\tilde{\sigma}\epsilon^j | \sigma \in G, 0 \leq j \leq w-1\}$, and the multiplication in $\widetilde{G}$ is given by $\widetilde{\sigma\tau} = \tilde{\sigma}\tilde{\tau}\epsilon^{\log_\gamma i(\sigma,\tau)}$, where $i(\sigma,\tau) = \frac{v_{\sigma\tau}}{v_\sigma\sigma(v_\tau)} \in \mathbb{F}_q^*$. For any $\tilde{\sigma} \in \widetilde{G}$, $\epsilon$ and $(\tilde{\sigma})^w$ are in the center of $\widetilde{G}$.*

*Proof.* By [Sect.5.1.2, 4], there exists such $v_\sigma \in K^*$ for $\sigma \in G$. The rest of the proof is trivial, which we refer to the proof of [Lemma 1, 2]. □

Let $M = P_1^{r_1}P_2^{r_2}\cdots P_n^{r_n}$ be the prime decomposition of $M$. We have the isomorphism:

$$G \cong (\mathbb{A}/M\mathbb{A})^* \cong (\mathbb{A}/P_1^{r_1}\mathbb{A})^* \times (\mathbb{A}/P_2^{r_2}\mathbb{A})^* \times \cdots \times (\mathbb{A}/P_n^{r_n}\mathbb{A})^*.$$

Different from the case of characteristic 0, now each $(\mathbb{A}/P_i^{r_i}\mathbb{A})^*$ is not always cyclic. But we have the decomposition $(\mathbb{A}/P_i^{r_i}\mathbb{A})^* \cong (\mathbb{A}/P_i^{r_i}\mathbb{A})^{(1)} \times (\mathbb{A}/P_i\mathbb{A})^*$, where $(\mathbb{A}/P_i^{r_i}\mathbb{A})^{(1)}$ is a $p$-group of order $|P_i|^{r_i-1}$ and $(\mathbb{A}/P_i\mathbb{A})^*$ is a cyclic group of order $|P_i|-1$, see [Prop.1.6, 5]. Let $\sigma_{P_i} \in G$ associate to a generator of $(\mathbb{A}/P_i\mathbb{A})^*$. Then we have

$$G = G^{(p)} \times G^{'},$$

where $G^{'} =< \sigma_{P_1}, \cdots, \sigma_{P_n} >=< \sigma_{P_1} > \times \cdots \times < \sigma_{P_n} >$, and $G^{(p)}$ is the $p$-Sylow subgroup of $G$. In fact, $G^{(p)} \cong (\mathbb{A}/P_1^{r_1}\mathbb{A})^{(1)} \times \cdots \times (\mathbb{A}/P_n^{r_n}\mathbb{A})^{(1)}$.

Let $\widetilde{G^{(p)}}$ and $\widetilde{G'}$ are the subgroups of $\widetilde{G}$ consisting of all liftings of the elements in $G^{(p)}$ and in $G^{'}$ respectively. It is easy to see they are normal subgroups of $\widetilde{G}$.

**Lemma 2.2.** *Let $\widetilde{G}^{(p)}$ be the p-Sylow subgroup of $\widetilde{G}$. Then*

$$\widetilde{G}^{(p)} \cong \widetilde{G^{(p)}}/ < \epsilon > \cong G^{(p)}.$$

*Furthermore, we have $\widetilde{G} = \widetilde{G}^{(p)} \times \widetilde{G'}$.*

*Proof.* Since $|\widetilde{G}| = w|G| = |G^{(p)}| \cdot |\widetilde{G'}|$, we have $|\widetilde{G}^{(p)}| = |G^{(p)}|$. Since each $\tilde{\sigma} \in \widetilde{G}^{(p)}$ is a lifting of some $\sigma \in G^{(p)}$, all liftings of $\sigma$ in $\widetilde{K}$ are $\tilde{\sigma}\epsilon^i (0 \leq i \leq w-1)$. Then the map $\sigma \mapsto \tilde{\sigma} \mod < \epsilon >$ gives the isomorphism $G^{(p)} \cong \widetilde{G^{(p)}}/ < \epsilon >$. Since the order of $\epsilon$ equals to $q-1$, we see $\tilde{\sigma}$ is the unique lifting of $\sigma$ belonging to $\widetilde{G}^{(p)}$. Thus $\sigma \mapsto \tilde{\sigma}$ gives the isomorphism $G^{(p)} \cong \widetilde{G}^{(p)}$.

Since $(\widetilde{G}^{(p)})^w = \widetilde{G}^{(p)}$, we have that $\widetilde{G}^{(p)}$ lies in the center of $\widetilde{G}$, and so it is a normal subgroup. In addition, as $|\widetilde{G}| = |\widetilde{G}^{(p)}| \cdot |\widetilde{G'}|$ and $(|\widetilde{G}^{(p)}|, |\widetilde{G'}|) = 1$, we have $\widetilde{G} = \widetilde{G}^{(p)} \times \widetilde{G'}$. □

Next we consider the group $\widetilde{G'}$. For the generators $\sigma_{P_i} \in G^{'}$ $(1 \leq i \leq n)$, we fix a lifting $\tilde{\sigma}_{P_i}$ of $\sigma_{P_i}$ in $\widetilde{K}$ as follows.

If $P_i \neq P, Q$, we define

$$\tilde{\sigma}_{P_i}(\sqrt[w]{u}) = \sqrt[w]{u}.$$

In fact, by [Sect.3.3, 4.3 and 5.1, 2], we have $\sigma_{P_i}(u) = u$.

If $P_i = P$ or $Q$, we define

$$\tilde{\sigma}_{P_i}(\sqrt[w]{u}) = v_{\sigma_{P_i}} \sqrt[w]{u},$$

where $v_{\sigma_{P_i}} \in K$ is given by

$$v_{\sigma_P} = \left( \sqrt[w]{(-1)^{d_Q}} \sin \mathbf{c}_{\sigma_P} \right)^{-1} \quad \text{and} \quad v_{\sigma_Q} = \left( \sqrt[w]{(-1)^{d_P}} \sin \mathbf{c}_{\sigma_Q} \right)^{-1},$$

here $\mathbf{c}_{\sigma_P}$ and $\mathbf{c}_{\sigma_Q}$ are defined in [Sect.4.2.5, 2]. By [Sect.3.4.2 and 3.4.3, 4], we have $\frac{u}{\sigma_P(u)} = (\sqrt[w]{(-1)^{d_Q}} \sin \mathbf{c}_{\sigma_P})^w$ with $\sqrt[w]{(-1)^{d_Q}} \sin \mathbf{c}_{\sigma_P} \in K^*$. Similarly for $\tilde{\sigma}_Q$.

Then we have

$$\widetilde{G'} =< \tilde{\sigma}_{P_1}, \cdots, \tilde{\sigma}_{P_n}, \epsilon > .$$

Now we study the relations among these generators of $\widetilde{G'}$. First $\epsilon$ commutes with each generator. By the calculations in [Sect.3.5, 3.6 and 5.1, 2], we see that the generators $\tilde{\sigma}_{P_i}$ commute with each other except for the relation

$$\tilde{\sigma}_P \tilde{\sigma}_Q = \tilde{\sigma}_Q \tilde{\sigma}_P \epsilon^{-1}.$$

In addition, by definition, if $P_i \neq P, Q$, then ord$(\tilde{\sigma}_{P_i})$=ord$(\sigma_{P_i})$. Finally, we need to compute the orders of $\tilde{\sigma}_P$ and $\tilde{\sigma}_Q$.

Let $L \in S_M$ and let $I_L$ be the inertia group of $L$ in $K$. It is known that $I_L \cong (\mathbb{A}/L^r\mathbb{A})^* \cong (\mathbb{A}/L^r\mathbb{A})^{(1)} \times (\mathbb{A}/L\mathbb{A})^*$, where $r$ is the maximal power of $L$ in $M$. We fix an inertia group $\widetilde{I}_L$ of $L$ in $\widetilde{K}$. Let $\widetilde{L}$ be a prime ideal in $\widetilde{K}$ over $L$ such that the inertia group $I(\widetilde{L}/L) = \widetilde{I}_L$. Let $\widetilde{G}_i$ be the $i$-th ramification group of $L|\widetilde{L}$, $i \geq -1$. Then by [III 8.6, 6], $\widetilde{G}_0 = \widetilde{I}_L$, $\widetilde{I}_L/\widetilde{G}_1$ is cyclic of order relatively prime to $p$, and $\widetilde{G}_1$ is the unique $p$-Sylow subgroup of $\widetilde{I}_L$, which is in the center of $\widetilde{I}_L$ by Lemma 2.1. Put $G_L =< \sigma_L >$ and $\widetilde{G}_L = \widetilde{G_L} \bigcap \widetilde{I}_L$, where $\widetilde{G_L}$ is the subgroup of $\widetilde{G}$ consisting of all liftings of the elements of $G_L$. Write $e_L$ the ramification index of $L$ in the extension $\widetilde{K}/K$.

**Proposition 2.3.** *We have $\widetilde{G}_1 \cong (\mathbb{A}/L^r\mathbb{A})^{(1)}$ and $\widetilde{G}_L$ is a cyclic group generated by some lifting of $\sigma_L$. Furthermore $\widetilde{I}_L = \widetilde{G}_1 \times \widetilde{G}_L$. In particular, all the liftings of $\sigma_L$ have the same order $e_L \cdot \mathrm{ord}(\sigma_L)$.*

*Proof.* Set $H =< \epsilon >$. The canonical homomorphism $\widetilde{I}_L \to I_L$, $\tilde{\sigma} \mapsto \tilde{\sigma}|_K$, induces an isomorphism $\widetilde{I}_L/(\widetilde{I}_L \cap H) \cong I_L$. Since $\widetilde{K}$ is abelian over $K$, we see $\widetilde{I}_L \cap H$ is the inertia group of $L$ in the field extension $\widetilde{K}/K$ by [Prop. 9.8, 5]. So the order of $\widetilde{I}_L \cap H$ is $e_L$. Notice that $\widetilde{G}_L \cap H = \widetilde{I}_L \cap H$. Hence $|\widetilde{I}_L| = e_L|I_L|$ and $|\widetilde{G}_L| = e_L|G_L|$.

Since $\widetilde{G}_1$ is in the center of $\widetilde{I}_L$ and $\widetilde{I}_L/\widetilde{G}_1$ is cyclic, we see $\widetilde{I}_L$ is abelian. Noting that $(|\widetilde{G}_L|, |\widetilde{G}_1|) = 1$ and $|\widetilde{I}_L| = |\widetilde{G}_1| \cdot |\widetilde{G}_L|$, we have $\widetilde{I}_L = \widetilde{G}_1 \times \widetilde{G}_L$. So $\widetilde{G}_L \cong \widetilde{I}_L/\widetilde{G}_1$ is cyclic. Since $I_L = \widetilde{I}_L|_K = \widetilde{G}_1|_K \times \widetilde{G}_L|_K = \widetilde{G}_1|_K \times G_L$ and $\widetilde{G}_1|_K \cong \widetilde{G}_1/(\widetilde{G}_1 \cap H) = \widetilde{G}_1$, we have $\widetilde{G}_1 \cong (\mathbb{A}/L^r\mathbb{A})^{(1)}$.

As $\widetilde{G}_L|_K = G_L$, there exists a lifting $\tilde{\sigma}_L$ of $\sigma_L$ belonging to $\widetilde{G}_L$. Since the order of $\epsilon$ is a factor of $|G_L|$, all the liftings of $\sigma_L$ have the same order. If the order of $\tilde{\sigma}_L$ is less than $|\widetilde{G}_L|$, then it is easy to show that the order of each element is also less than $|\widetilde{G}_L|$. But $\widetilde{G}_L$ is cyclic. Hence $\tilde{\sigma}_L$ is a generator of $\widetilde{G}_L$. □

If $P_i \neq P$ and $Q$, then $e_{P_i} = 1$. Now we compute the ramification indices $e_P$ and $e_Q$ of $P$ and $Q$ in the extension $\widetilde{K}/K$.

Let $R$ be a monic irreducible polynomial in $\mathbb{A}$ and $A \in \mathbb{A}$ be coprime to $R$. Recall that the $(q-1)$th residue symbol $(\frac{A}{R}) \in \mathbb{F}_q^*$ is defined by

$$(\frac{A}{R}) \equiv A^{\frac{|R|-1}{q-1}} \mod R.$$

Let $v_P$ be the normalized valuation in $k^{ab}$ associated to $P$ defined in [Sect.6, 2]. By [Prop.6.2, 2] we have

$$v_P(\mathbf{e}(\mathbf{a}_{PQ})) \equiv \log_\gamma(\frac{Q}{P}) \quad \text{and} \quad v_Q(\mathbf{e}(\mathbf{a}_{PQ})) \equiv -\log_\gamma(\frac{P}{Q}) \pmod w.$$

and $v_P(P)$ equals to the ramification index $|P|^{r-1}(|P|-1)$ of $P$ in $K/k$, where $r$ is the maximal power of $P$ in $M$. Thus $v_P(\sqrt{P}) \equiv \frac{w}{2}d_P \pmod w$. Similarly, we have $v_Q(\sqrt{Q}) \equiv \frac{w}{2}d_Q \pmod w$.

Furthermore, by the reciprocity law $(\frac{Q}{P}) = (-1)^{d_P d_Q}(\frac{P}{Q})$, see [Theorem 3.5, 5], we have

$$v_P(u) \equiv \log_\gamma(\frac{P}{Q}) \quad \text{and} \quad v_Q(u) \equiv -\log_\gamma(\frac{Q}{P}) \pmod w,$$

By [III 7.3, 6], we have $e_P = \frac{w}{(w, v_P(u))}$ and $e_Q = \frac{w}{(w, v_Q(u))}$, so

$$e_P = \frac{w}{(w, \log_\gamma(\frac{P}{Q}))} \quad \text{and} \quad e_Q = \frac{w}{(w, \log_\gamma(\frac{Q}{P}))}.$$

In particularly, if $2|d_P d_Q$, we have $e_P = e_Q$. Finally we get the following theorem.

**Theorem 2.4.** *We have $\widetilde{G} = \widetilde{G}^{(p)} \times \widetilde{G'}$, where $\widetilde{G}^{(p)}$ is the p-Sylow subgroup of $\widetilde{G}$, and $\widetilde{G'} =< \tilde{\sigma}_{P_1}, \cdots, \tilde{\sigma}_{P_n}, \epsilon >$. The generators $\tilde{\sigma}_{P_i}$ and $\epsilon$ commute with each other except for the relation $\tilde{\sigma}_P \tilde{\sigma}_Q = \tilde{\sigma}_Q \tilde{\sigma}_P \epsilon^{-1}$. In addition, for $P_i \neq P, Q$, we have* $\mathrm{ord}(\tilde{\sigma}_{P_i})$=$\mathrm{ord}(\sigma_{P_i})$, *and for $P_i = P$ or $Q$, we have*

$$\mathrm{ord}(\tilde{\sigma}_P) = \frac{w}{(w, \log_\gamma(\frac{P}{Q}))} \cdot \mathrm{ord}(\sigma_P) \quad \textit{and} \quad \mathrm{ord}(\tilde{\sigma}_Q) = \frac{w}{(w, \log_\gamma(\frac{Q}{P}))} \cdot \mathrm{ord}(\sigma_Q).$$

## 3. The genus formula

In this section we compute the genus of $\widetilde{K}$. We calculate it by using Hasse's genus formula on Kummer extensions, which states that for a $m$-th Kummer extension $E/F$ of algebraic function fields, $m$ is relatively prime to the characteristic of $\mathbb{F}_F$, $[E:F] = m$, and there exists $a \in F$ such that $E = F(\sqrt[m]{a})$, then we have

$$g_E = 1 + \frac{m}{[\mathbb{F}_E : \mathbb{F}_F]} \left[ g_F - 1 + \frac{1}{2} \sum_{\mathfrak{p} \in \mathbb{P}_F} \left(1 - \frac{1}{e_\mathfrak{p}}\right) \deg \mathfrak{p} \right],$$

where $g_E$ and $g_F$ are the genus of $E$ and $F$ respectively, $\mathbb{F}_E$ and $\mathbb{F}_F$ are the constant fields of $E$ and $F$ respectively, $\mathbb{P}_F$ is the set of primes of $F$, and $e_\mathfrak{p}$ is the ramification index of $\mathfrak{p}$ in $E/F$. See [III 7.3, 6].

We first need to compute the genus of $K$. In [3], Hayes calculated it in the case that the conductor $M$ of $K$ has only one prime factor. Here we use the Riemann-Hurwitz theorem to calculate the genus of $K$ in general case.

Let $E/F$ be a finite, separable, geometric extension of algebraic function fields, the Riemann-Hurwitz theorem says

$$2g_E - 2 = [E:F](2g_F - 2) + \deg D_{E/F},$$

where $D_{E/F}$ is the different divisor of $E/F$, see [Theorem 7.16, 5].

Recall that $M$ has the prime decomposition $M = P_1^{r_1} P_2^{r_2} \cdots P_n^{r_n}$.

**Proposition 3.1.** *We have*

$$g_K = \left[ \frac{q-2}{2(q-1)} - 1 \right] \Phi(M) + \frac{1}{2} \sum_{i=1}^{n} s_i d_i \Phi(M/P_i^{r_i}) + 1,$$

*where $d_i = \deg P_i$, $s_i = r_i \Phi(P_i^{r_i}) - q^{d_i(r_i - 1)}$ and $\Phi(M) = |(\mathbb{A}/M)^*|$.*

*Proof.* Let $L \in S_M$ and $r$ the maximal power of $L$ in $M$. Put $d = \deg L$ and $E = k(\mathbf{e}_C(\frac{\bar{\pi}}{L^r}))$. Let $i_{K/E}$ be the homomorphism from the divisor group of $E$ to that of $K$ defined by $i_{K/E}(\mathfrak{p}) = \sum_{\mathfrak{P}|\mathfrak{p}} e(\mathfrak{P}/\mathfrak{p})\mathfrak{P}$, where $\mathfrak{p}$ is a prime of $E$. By [Page 87, 5], we have

$$D_{K/k} = D_{K/E} + i_{K/E}(D_{E/k}).$$

It is known that $L$ is totally ramified in $E/k$. Let $\mathfrak{l}$ be the unique prime ideal of $E$ over $L$. Let $\mathfrak{L}_1, \mathfrak{L}_2, \cdots, \mathfrak{L}_g$ be all the different prime ideals of $K$ over $L$. For each $\mathfrak{L}_j (1 \leq j \leq g)$, we now calculate its coefficient in $D_{K/k}$. Since $\mathfrak{L}_j$ is unramified in

$K/E$, its coefficient in $D_{K/E}$ is 0. By [Theorem 4.1, 3], the coefficient of $\mathfrak{l}$ in $D_{E/k}$ is $s = r\Phi(L^r) - q^{d(r-1)}$, and so

$$i_{K/E}(s\mathfrak{l}) = s i_{K/E}(\mathfrak{l}) = s \sum_{\mathfrak{L}_j|\mathfrak{l}} e(\mathfrak{L}_j/\mathfrak{l})\mathfrak{L}_j = s \sum_{\mathfrak{L}_j|\mathfrak{l}} \mathfrak{L}_j.$$

Thus for each $\mathfrak{L}_j$, its coefficient in $D_{K/k}$ is $s$.

Let $\infty$ be the infinite prime of $k$. There are $\frac{\Phi(M)}{q-1}$ different infinite primes in $K$ over $\infty$, and the ramification index is $q-1$. Thus the coefficient of each infinite prime of $K$ in $D_{K/k}$ is $q-2$, also see Corollary in Page 85 in [5].

We have determined $D_{K/k}$. By Riemann-Hurwitz theorem, we get the desired formula. □

The constant field of $\widetilde{K}$ is clearly $\mathbb{F}_q$. In last section we have computed the ramification indices in $\widetilde{K}/K$ of all the finite primes of $K$. To calculate the genus of $\widetilde{K}$, we need to compute that of the infinite primes.

Let $k_\infty \subset \Omega$ be the completion of $k$ at the place $1/T$. Let $K^+ = K \cap k_\infty$ be the maximal real subfield of $K$. By [Sect.4.3, 2], we know $\sin\mathbf{a}_{PQ} = \mathbf{e}(\mathbf{a}_{PQ}) \in K^+$. It is known that for any monic square-free polynomial $f(T)$ in $\mathbb{F}_q[T]$ with even degree, we have $\sqrt{f(T)} \in k_\infty$. So $u \in K^+$.

Let $E = K^+(\sqrt[w]{u})$. Then $\widetilde{K} = EK$ and $[E : K^+] = w$. Let $\infty$ be any infinite prime of $K^+$, $\infty_1$ an infinite prime of $K$ over $\infty$, $\infty_2$ an infinite prime of $E$ over $\infty$, and $\widetilde{\infty}$ an infinite prime of $\widetilde{K}$ over $\infty_1$. By [Theorem 12.14, 5], the ramification index $e(\infty_1/\infty) = w$. Then by Abhyankar's Lemma, see [III 8.9, 6], the ramification index $e(\widetilde{\infty}/\infty) = w$. Since $e(\widetilde{\infty}/\infty) = e(\widetilde{\infty}/\infty_1) \cdot e(\infty_1/\infty)$, we have $e(\widetilde{\infty}/\infty_1) = 1$. Thus $\infty_1$ is unramified in $\widetilde{K}/K$. Since $\widetilde{K}$ is Galois over $K$, all infinite primes of $K$ are unramified in $\widetilde{K}/K$. Now we can get the genus formula of $\widetilde{K}$.

**Theorem 3.2.** *We have*

$$g_{\widetilde{K}} = 1 + w\left[g_K - 1 + \frac{1}{2}\left(1 - \frac{1}{e_P}\right) d_P \Phi(M/P^{r_P}) + \frac{1}{2}\left(1 - \frac{1}{e_Q}\right) d_Q \Phi(M/Q^{r_Q})\right],$$

*where $r_P$ and $r_Q$ are the maximal powers of $P$ and $Q$ in $M$ respectively, $e_P = \frac{w}{(w, \log_\gamma(\frac{P}{Q}))}$ and $e_Q = \frac{w}{(w, \log_\gamma(\frac{Q}{P}))}$.*

*Proof.* By Hasse's formula, we have

$$g_{\widetilde{K}} = 1 + w\left[g_K - 1 + \frac{1}{2} \sum_{\substack{\text{prime } \mathfrak{p} \text{ in } K \\ \mathfrak{p}|P \text{ or } \mathfrak{p}|Q}} \left(1 - \frac{1}{e_{\mathfrak{p}}}\right) \deg\mathfrak{p}\right],$$

where the sum is over all the uneqvivalent primes over $P$ or $Q$, $e_{\mathfrak{p}}$ is the ramification index of $\mathfrak{p}$ in $\widetilde{K}/K$, and $\deg\mathfrak{p} = f(\mathfrak{p}/P)d_P$.

We assume that there are $g_P$ and $g_Q$ different prime ideals in $K$ over $P$ and $Q$ respectively. Then

$$g_{\widetilde{K}} = 1 + w\left[g_K - 1 + \frac{1}{2}\left(1 - \frac{1}{e_P}\right) g_P f_P d_P + \frac{1}{2}\left(1 - \frac{1}{e_Q}\right) g_Q f_Q d_Q\right],$$

where $f_P$ and $f_Q$ are the residue class degrees of $P$ and $Q$ in $K/k$ respectively.

Since $g_P f_P = \Phi(M/P^{r_P})$ and $g_Q f_Q = \Phi(M/Q^{r_Q})$, we get the formula. □

## 4. The general case

In this section, we consider the general case. We follow the method in [7] to determine the Galois group in this case.

Let $\widetilde{K} = K(\sqrt[w]{u})$, where $u = u_{P_1Q_1} \cdots u_{P_mQ_m}$ and $P_i < Q_i \in S_M$. We assume $(P_i, Q_i) \neq (P_j, Q_j)$ for $i \neq j$. Then $\widetilde{K}$ is also a quasi-cyclotomic function field over $k$, whose degree over $K$ equals to $q-1$. Notice that there may exist $P_i$ or $Q_i$ appearing repeatedly in this sequence $P_1, Q_1, \cdots, P_m, Q_m$.

Let $\widetilde{G} = \mathrm{Gal}(\widetilde{K}/k)$. Applying the same arguments, Lemma 2.1 and Lemma 2.2 are also true here. Then we have $\widetilde{G} = \widetilde{G}^{(p)} \times \widetilde{G'}$, where the meanings of the notations are similar as before.

We consider the field $\widehat{K} = K(\sqrt[w]{u_{P_1Q_1}}, \cdots, \sqrt[w]{u_{P_mQ_m}})$. It is Galois over $k$ with degree $w^m\Phi(M)$. Each element of $G$ has $w^m$ liftings to $\widehat{K}$. For each $P \in S_M$, we define a lifting $\hat{\sigma}_P$ of $\sigma_P$ to $\widehat{K}$ as before. If $P \neq P_i, Q_i$, then $\hat{\sigma}_P(\sqrt[w]{u_{P_iQ_i}}) = \sqrt[w]{u_{P_iQ_i}}$. Otherwise, $\hat{\sigma}_{P_i}(\sqrt[w]{u_{P_iQ_i}}) = v_{\sigma_{P_i}} \sqrt[w]{u_{P_iQ_i}}$ and $\hat{\sigma}_{Q_i}(\sqrt[w]{u_{P_iQ_i}}) = v_{\sigma_{Q_i}} \sqrt[w]{u_{P_iQ_i}}$. We denote the restriction of $\hat{\sigma}_P$ on $\widetilde{K}$ by $\tilde{\sigma}_P$. Then the group $\widetilde{G'}$ is generated by $\left\{\tilde{\sigma}_P, \epsilon | P \in S_M\right\}$, where $\epsilon$ is defined as before.

Now we consider the relations between these generators. If $P \in S_M$ and $P \neq P_i, Q_i (1 \leq i \leq m)$, it is easy to see $\tilde{\sigma}_P$ commutes with all generators, and the order of $\tilde{\sigma}_P$ equals to that of $\sigma_P$. By the discussion in section 2, we have $\hat{\sigma}_{P_i}\hat{\sigma}_{Q_i} = \hat{\sigma}_{P_i}\hat{\sigma}_{Q_i}\epsilon_i^{-1}$ for each $i$, where $\epsilon_i$ acts trivially on $K$ and $\sqrt[w]{u_{P_jQ_j}}$ for $j \neq i$, and $\epsilon_i(\sqrt[w]{u_{P_iQ_i}}) = \gamma \sqrt[w]{u_{P_iQ_i}}$. Since $\epsilon_i|_{\widetilde{K}} = \epsilon$, we see $\tilde{\sigma}_{P_i}\tilde{\sigma}_{Q_i} = \tilde{\sigma}_{Q_i}\tilde{\sigma}_{P_i}\epsilon^{-1}$. The other multiplication relations among $\tilde{\sigma}_{P_1}, \tilde{\sigma}_{Q_1}, \cdots, \tilde{\sigma}_{P_m}, \tilde{\sigma}_{Q_m}$ are complicated, but it is easy to determine. Since Proposition 2.3 is also true here, the orders of $\tilde{\sigma}_{P_i}$ and $\tilde{\sigma}_{Q_i}$ can be determined by the ramification natures of $P_i$ and $Q_i$ in $\widetilde{K}/K$. In other words, let $P$ be one of $P_1, Q_1, \cdots, P_m, Q_m$, let $L_1, \cdots, L_s$ be all the monic prime factors paired with $P$ with the form $u_{PL_i} (1 \leq i \leq s)$, and let $R_1, \cdots, R_t$ be all the monic prime factors paired with $P$ with the form $u_{R_iP} (1 \leq i \leq t)$, then $\mathrm{ord}(\tilde{\sigma}_P) = \frac{w}{\left(w, \log_\gamma(\frac{P}{L_1 \cdots L_s}) - \log_\gamma(\frac{P}{R_1 \cdots R_t})\right)} \cdot \mathrm{ord}(\sigma_P)$. Therefore, we determine the group $\widetilde{G}$ by generators and relations.

In fact, we have got the ramification indexes $e_{P_i}, e_{Q_i} (1 \leq i \leq m)$ in the above paragraph. We can also get that all the infinite primes of $K$ are unramified in the field extension $\widetilde{K}/K$ by applying the same argument as the paragraph above Theorem 3.2. Hence, we can get the genus formula in the general case by applying the same argument as Theorem 3.2.

## References

[1] S. Bae, E.-U. Gekeler, P. Kang and L. Yin, *Anderson's double complex and gamma monomials for rational function fields*, Trans. Amer. Math. Soc. **355** (2003), 3463-3474.

[2] S. Bae, L. Yin, *Carlitz-Hayes plus Anderson's epsilon*, J. reine angew. Math. **571** (2004), 19-37.

[3] D.R. Hayes, *Explicit class field theory for rational function fields*, Trans. Amer. Math. Soc. **189** (1974), 77-91.

[4] C. Liu, L. Yin, *Double coverings for quadratic extensions and function fields*, J. Number Theory **130** (2010), 469-477.

[5] M. Rosen, *Number theory in function fields*, GTM 210, Springer-Verlag, 2002.

[6] H. Stichtenoth, *Algebraic function fields and codes*, Springer-Verlag, 1993.

[7] L. Yin, C. Zhang, *Arithmetic of quasi-cyclotomic fields*, J. Number Theory **128** (2008), 1717-1730.

Department of Mathematical Sciences, Tsinghua University, Beijing 100084, P.R. China
*E-mail address*: `sham07@mails.tsinghua.edu.cn`

Department of Mathematical Sciences, Tsinghua University, Beijing 100084, P.R. China
*E-mail address*: `lsyin@math.tsinghua.edu.cn`